

Heat Emergencies	
ADULT	PEDIATRIC
BLS Procedures	
<p>Place patient in cool, shaded environment Initiate active cooling measures</p> <ul style="list-style-type: none"> • Remove clothing • Fan the patient, or turn on air conditioner • Apply ice packs to axilla, groin, back of neck • Other active cooling measures as available <p>Administer oxygen as indicated If patient is altered, determine blood glucose level If less than 60 mg/dl refer to Policy 705.03</p>	
ALS Standing Orders	
<p>If patient is altered, determine blood glucose if not already performed by BLS personnel or post oral glucose administration</p> <p>If less than 60 mg/dl, refer to Policy 705.03</p> <p>IV/IO access</p> <p>Normal Saline</p> <ul style="list-style-type: none"> • IV/IO bolus – 1 Liter <ul style="list-style-type: none"> ○ Caution with cardiac and/or renal history ○ Repeat x 1 for persistent hypotension 	<p>If patient is altered, determine blood glucose if not already performed by BLS personnel or post oral glucose administration</p> <p>If less than 60 mg/dl, refer to Policy 705.03</p> <p>IV/IO access</p> <p>Normal Saline</p> <ul style="list-style-type: none"> • IV/IO bolus – 20 mL/kg <ul style="list-style-type: none"> ○ Caution with cardiac and/or renal history ○ Repeat x 1 for persistent hypotension
Base Hospital Orders only	
Consult with ED Physician for further treatment measures	